arXiv:1010.6119v1 [math.AG] 29 Oct 2010

A q-analog of the Fibonacci numbers via complete quadrics

Mahir Bilen Can
Micheal Joyce

November 1, 2010

Abstract

We analyze the fixed variety Y_n of a special unipotent element acting on the space of $(n-1)$-dimensional complete quadrics. We find a cell decomposition of Y_n, and compute its Poincaré polynomial. The cells are indexed by compositions of n with odd parts and the Poincaré polynomial of Y_n is a q-analog of the n^{th} Fibonacci number. We characterize the containment relations of the cell closures combinatorially and find a recurrence for the number of irreducible components of Y_n.

1 Introduction

Let $\mathscr{B}$ be the variety of n-step flags in $\mathbb{C}^n$:

$$\mathcal{F} : 0 = V_0 \subset V_1 \subset \cdots \subset V_{n-1} \subset V_n = \mathbb{C}^n,$$

where $\dim V_i = i$ for $i = 0, \dots, n$. Let u be a unipotent element from the special linear group $SL_n(\mathbb{C})$. The variety of fixed points $\mathscr{B}^u := \{\mathcal{F} \in \mathscr{B} : \ u \cdot \mathcal{F} = \mathcal{F}\}$ has an intricate geometry and a rich combinatorial structure.

It was shown by Springer ([10], [11]) that for each $i \geq 0$, the cohomology space $H^i(\mathscr{B}^u, \mathbb{Q})$ affords a representation of the symmetric group S_n, and the top non-vanishing cohomology space is an irreducible S_n-module. Spaltenstein [9] proved that all irreducible components of $\mathscr{B}^u$ have the same dimension $e(u)$ which is equal to the half of the dimension of the top non-vanishing cohomology space $H^{2e(u)}(\mathscr{B}^u, \mathbb{Q})$.

The flag variety $\mathcal{B}$ is an example of a wonderful SL_n-variety. Another frequently studied wonderful SL_n-variety is X_n, the space of complete quadrics in an n-dimensional affine space ([5], [6]). A complete quadric can be defined inductively by giving a quadric hypersurface Q in the affine space and a complete quadric in the singular locus of Q, which is itself an affine space.

We investigate the analog of the Springer fiber for X_n,

$$Y_n := X_n^u = \{x \in X_n : \ u \cdot x = x\},$$

in the special case where u has Jordan type (n), i.e. consists of a single Jordan block. Y_n has a cell decomposition and the cell containment relations can be described combinatorially in terms of compositions of n whose parts are all odd. In fact, the cells form a graded poset and a meet semilattice.

The Poincaré polynomial of Y_n turns out to be a q-analog of the Fibonacci numbers:

$$P_{Y_n}(q) = \sum_{i=0}^{n} \binom{n-i}{i} q^i,$$

satisfying the recurrence

$$P_{Y_n}(q) = P_{Y_{n-1}}(q) + qP_{Y_{n-2}}(q), \ \ P_{Y_0}(q) = P_{Y_1}(q) = 1.$$

The number a_n of irreducible components of Y_n satisfies the recurrence: $a_n = a_{n-1} + a_{n-3}$. Unlike the case of Springer fibers for $\mathcal{B}$, the irreducible components of Y_n are never equidimensional when $n \geq 5$.

2 Preliminaries

2.1 Compositions with Odd Parts

A *composition* of $n \in \mathbb{N}$ is an ordered sequence of positive integers (called parts) that sum to n. Let F_n denote the set of compositions of n whose parts are odd. It is well known that $|F_n|$ is given by the n^{th} Fibonacci number (c.f. [12, Exercise I.14(d)]).

We partition F_n into a disjoint union of two subsets:

$$F_n = F_n' \sqcup F_n'',$$

where F'_n, respectively F''_n, consists of compositions $\gamma = (\gamma_1, \gamma_2, \ldots, \gamma_k) \in F_n$ with $\gamma_1 = 1$, respectively $\gamma_1 > 1$. There are bijections $\phi : F'_n \to F_{n-1}$,

$$\phi(\gamma) = (\gamma_2, \gamma_3, \ldots, \gamma_k) \tag{2.1}$$

and $\psi : F''_n \to F_{n-2}$,

$$\psi(\gamma) = (\gamma_1 - 2, \gamma_2, \gamma_3, \ldots, \gamma_k). \tag{2.2}$$

2.2 Generalities on Quadrics

We fix an algebraically closed field $\mathbb{K}$ of characteristic 0 and let V denote a $\mathbb{K}$-vector space of dimension n. A quadric hypersurface in V is the locus of $v \in V$ such that $q(v) = 0$ for some non-zero quadratic form q on V. Two such quadratic forms produce the same quadric hypersurface if and only if one is a scalar multiple of the other. Via the polarization trick, quadratic forms are equivalent to symmetric bilinear forms. Moreover, if we fix an ordered basis

$$\mathcal{E} = \{e_1, e_2, \ldots, e_n\}, \tag{2.3}$$

then a symmetric bilinear form b is given by a symmetric matrix $A = \big(b(e_i, e_j)\big)$. Let $\mathrm{Sym}_n = \mathrm{Sym}_n(\mathbb{K})$ denote the vector space of symmetric n-by-n matrices with entries in $\mathbb{K}$. The space of all quadric hypersurfaces can therefore be identified with the projectivization, $\mathbb{P}(\mathrm{Sym}_n)$; it has dimension $\frac{n(n+1)}{2} - 1$.

2.3 Complete Quadrics

A quadric hypersurface is non-degenerate if its defining quadratic form q is non-degenerate; equivalently, if its defining symmetric matrix A is invertible. Let $\mathcal{Q} = \mathcal{Q}_n$ denote the open subset of $\mathbb{P}(\mathrm{Sym}_n)$ corresponding to non degenerate quadrics. The space X_n of *complete quadrics* is an alternative compactification of $\mathcal{Q}$ of classical origin, going back to Chasles [4] and Schubert [8]. A complete quadric in V is a pair $(\mathcal{F}, Q) = (\mathcal{F}, \{Q_1, Q_2, \ldots, Q_k\})$, where

$$\mathcal{F} : 0 = V_0 \subset V_1 \subset V_2 \subset \cdots \subset V_k = V$$

is a partial flag of subspaces in V, and Q_i, for $i = 1, \ldots, k$, is a non-degenerate quadric in V_i/V_{i-1}. Equivalently, a a complete quadric can be described by a pair $(\mathcal{F}, \widetilde{Q}) = (\mathcal{F}, \{\widetilde{Q_1}, \widetilde{Q_2}, \ldots, \widetilde{Q_k}\})$ where $\mathcal{F}$ is as above and $\widetilde{Q_i}$ is a quadric defined on V_i whose singular locus is V_{i-1}. Note that a non-degenerate quadric is a complete quadric with $k = 1$.

X_n is a smooth projective variety and Q embeds into X_n as a dense open subset. See [5] and [6] for a more thorough description of X_n, including its construction as a wonderful SL_n-variety. The Poincaré polynomial of X_n is computed explicitly in [13] using a Bialynicki-Birula decomposition, and in [7] by point counting over finite fields.

Let $\mathcal{F}$ be a flag in $V = \mathbb{K}^n$ given by

$$0 = V_0 \subset V_1 \subset \cdots \subset V_k = V \tag{2.4}$$

and let $\gamma_i = \dim(V_i/V_{i-1})$. Note that $(\gamma_1, \gamma_2, \dots, \gamma_k)$ is a composition of n, called the *flag type* of $\mathcal{F}$. We say that our choice of ordered basis $\mathcal{E}$ in (2.3) is consistent with the flag $\mathcal{F}$ in the sense that the first $\gamma_1 + \gamma_2 + \cdots + \gamma_i$ vectors in $\mathcal{E}$ form a basis for V_i.

There is a natural right action of $SL_n = SL_n(\mathbb{K})$ on X_n. If $g \in SL_n$ and $\mathcal{F}$ is given by (2.4), $\mathcal{F} \cdot g$ is the flag

$$0 = V_0 \subset g^{-1}(V_1) \subset \cdots \subset g^{-1}(V_k) = V.$$

Note that g restricts to linear isomorphisms

$$g_i : g^{-1}(V_i)/g^{-1}(V_{i-1}) \longrightarrow V_i/V_{i-1} \tag{2.5}$$

The action on the non-degenerate quadrics defining the complete quadric is $Q \cdot g = (Q_1, Q_2, \dots, Q_k) \cdot g = (Q_1 \cdot g_1, Q_2 \cdot g_2, \dots Q_k \cdot g_k)$ is given by

$$Q_i \cdot g = g_i^{-1}(Q_i).$$

In the sequel, we will only need to consider the case when the action fixes the flag. In that case, let $\mathcal{E}_i$ the be the subset of $\mathcal{E}$ consisting of vectors in V_i but not in V_{i-1} (with the induced ordering). Then $\mathcal{E}_i$ is an ordered basis for V_i/V_{i-1}. Let A_i be an invertible symmetric matrix that gives the non-degenerate quadric hypersurface Q_i with respect to the basis $\mathcal{E}_i$. For $g \in SL_n$, use the composition γ to form a block decomposition of g. Then the matrix representing g_i (2.5) with respect to the basis $\mathcal{E}_i$ is simply the i^{th} diagonal block of g with respect to the γ-block decomposition. An invertible symmetric matrix that represents the non-degenerate quadric $Q_i \cdot g$ is given by $g_i A_i g_i^{\mathsf{T}}$. (The use of the given right action instead of the natural left action is dictated by this matrix formula for the action, as well as to be able to directly apply results from [2].)

2.4 Generalities on Posets

Definition 2.6. A poset P is *graded* if there exists a rank function $\ell : P \to \mathbb{N}$ such that

1. if $x < y$, then $\ell(x) < \ell(y)$;
2. if y covers x, then $\ell(y) = \ell(x) + 1$.

Note that Definition 2.6 differs from that in [12] in that we do not require P to have a unique minimal element $\hat{0}$ or to have a unique maximal element $\hat{1}$.

Definition 2.7. A poset P is called a *meet semi-lattice* if for every $a, b \in P$ there exists unique $a \wedge b \in P$ such that

1. $a \wedge b \leq a$ and $a \wedge b \leq b$;
2. if $c \leq a$ and $c \leq b$ for some $c \in P$, then $c \leq a \wedge b$.

3 Invariant Non-degenerate Quadrics

We recall results of [2] on unipotent invariant non-degenerate quadrics.

Proposition 3.1. *[2, Proposition 3.4] Let N be a nilpotent matrix and let $u = \exp(N)$. Let Q be a non-degenerate quadric in $V = \mathbb{K}^n$ defined by a symmetric matrix A. Then Q is fixed by u if and only if $N^{\mathsf{T}}A + AN = 0$.*

Given an arbitrary nilpotent transformation $\mathcal{N}$ of V whose Jordan type is the partition (n) having only one part, there is an isomorphism $V \cong \mathbb{K}^n$ such that $\mathcal{N}$ is given in matrix form by

$$N = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix}. \tag{3.2}$$

The associated unipotent transformation $\mathfrak{u} = \exp(\mathcal{N})$ is then given by the unipotent matrix $u = \exp(N)$.

Proposition 3.3. *[2, Corollaries 4.5, 4.6] Let $u = \exp(N)$, with N the n-by-n matrix given by (3.2).*

1. *If n is even, there are no non-degenerate quadrics fixed by u.*
2. *If $n = 2k + 1$ is odd, then the space Q^u of non-degenerate quadrics fixed by u is isomorphic to $\mathbb{A}^k$. Explicity, the symmetric matrices A corresponding to such quadrics (defined only up to scalar multiplication) are given by*

$$A = \begin{bmatrix} a_1 & 0 & a_2 & \cdots & a_{k+1} \\ 0 & -a_2 & 0 & \cdots & 0 \\ a_2 & 0 & a_3 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{k+1} & 0 & 0 & \cdots & 0 \end{bmatrix}. \tag{3.4}$$

Let T be the maximal torus of $SL(\mathbb{K}$ consisting of invertible diagonal matrices. We now determine the maximal subtorus T' of T that fixes Q^u.

Lemma 3.5. *A torus element $t \in T$ fixes Q^u if and only if t has the form*

$$\begin{pmatrix} \alpha^k & 0 & 0 & \cdots & 0 \\ 0 & \mu\alpha^{k-1} & 0 & \cdots & 0 \\ 0 & 0 & \mu^2\alpha^{k-2} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \mu^{2k}\alpha^{-k} \end{pmatrix},$$

where $\alpha \in \mathbb{K}^*$ and $\mu = \pm 1$.

In particular, T' consisting of

$$\lambda(t) = \begin{pmatrix} t^k & 0 & 0 & \cdots & 0 \\ 0 & t^{k-1} & 0 & \cdots & 0 \\ 0 & 0 & t^{k-2} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & t^{-k} \end{pmatrix} \tag{3.6}$$

for $t \in \mathbb{K}^*$ is a maximal subtorus of T that fixes Q^u.

Remark 3.7. The action of T' on Q^u has a unique fixed point given by

$$A_{(n)} = \begin{bmatrix} 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & -1 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & -1 & \cdots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \end{bmatrix}. \tag{3.8}$$

4 Invariant Complete Quadrics

Proposition 4.1. *Let* $V = \mathbb{K}^n$ *with standard basis* $\{e_1, e_2, \dots, e_n\}$, *let* N *be given by (3.2) and let* $u = \exp(N)$. *For each composition* γ *of* n, *there is a unique flag of type* γ *that is fixed by* u, *namely the sub-flag of type* γ *of the complete standard flag*

$$0 \subset \mathbb{K}e_1 \subset \mathbb{K}e_1 \oplus \mathbb{K}e_2 \subset \cdots \subset \mathbb{K}e_1 \oplus \mathbb{K}e_2 \oplus \cdots \oplus \mathbb{K}e_n = V.$$

Proof. We prove this by induction on the number k of parts of the composition γ. Observe that a flag

$$\mathcal{F} : 0 \subset V_1 \subset V_2 \subset \cdots \subset V_k = K^n$$

is fixed by u if and only if it is fixed by N, so assume $\mathcal{F}$ is N-fixed.

Let v be any non-zero vector in V_1 and let j be the largest integer such that v contains a non-zero e_j component. Then $v, N(v), N^2(v), \dots, N^{k-1}(v)$ are j linearly independent vectors in V_1. Since $\dim(V_1) = \gamma_1$, this implies $j \leq \gamma_1$. Thus $V_1 = \mathbb{K}e_1 \oplus \mathbb{K}e_2 \oplus \cdots \oplus \mathbb{K}e_{\gamma_1}$. Now apply the inductive hypothesis to the induced flag obtained by quotienting $\mathcal{F}$ by V_1, noting that the induced nilpotent transformation on V/V_1 has Jordan type $(n - \gamma_1)$ and, consequently, still has the form given by (3.2). □

Definition 4.2. For $\gamma \in F_n$, $X_{n,\gamma}$ is the subvariety of X_n consisting of complete quadrics whose flag has type γ. For a unipotent $u \in SL_n(\mathbb{K})$, define

$$Y_{n,\gamma} = X^u_{n,\gamma} = \{P = (\mathcal{F}, Q) \in X_{n,\gamma} : P \cdot u = P\}.$$

Observe that if

$$\mathcal{F} : 0 = V_0 \subset V_1 \subset \cdots \subset V_k = V$$

is the u-fixed flag of type $\gamma = (\gamma_1, \gamma_2, \dots, \gamma_k)$, then it is also N-fixed and the restriction of N to V_i/V_{i-1} is given by a γ_i-by-γ_i matrix of the form (3.2). Therefore, we may apply the results of Section 3 to study the non-degenerate u-fixed quadrics on V_i/V_{i-1}.

By part 1 of Proposition 3.3, there are no u-fixed quadrics of type γ if any of the γ_i are even. On the other hand, if each γ_i is odd, then applying part 2 of Proposition 3.3 shows that the space of non-degenerate u-fixed quadrics on V_i/V_{i-1} is an affine space of dimension $(\gamma_i - 1)/2$. By Remark 3.7, each of these affine spaces has a unique T' fixed point given by (3.8).

We summarize these observations in the following proposition.

Proposition 4.3. *Let* $\gamma = (\gamma_1, \ldots, \gamma_k) \in F_n$ *and let* $u \in SL_n(\mathbb{K})$ *be unipotent. Then,*

$$Y_{n,\gamma} \cong \mathbb{A}^{(\gamma_1-1)/2} \times \cdots \times \mathbb{A}^{(\gamma_k-1)/2} \cong \mathbb{A}^{(n-k)/2}.$$

Furthermore, there is a unique T'*-fixed point* A_γ *in* $Y_{n,\gamma}$.

5 Carrell and Goresky's *BB*-Theorem

In the sequel, let $\mathbb{K} = \mathbb{C}$.

Lemma 5.1. *Let* A_γ *be as in Proposition 4.3. Then,*

$$Y_{n,\gamma} = \{Q \in Y_n : \lim_{t\to 0} Q \cdot \lambda(t) = A_\gamma\}.$$

Proof. Since each $Y_{n,\gamma}$ has a product structure, it suffices to check the result for the case $\gamma = (n)$ for n odd, i.e. when the flag $\mathcal{F}$ is simply

$$\mathcal{F} : 0 \subset V.$$

The action of $\lambda(t)$ given by (3.6) on a non-degenerate quadric in Y_n given by A in (3.4) is

$$A \cdot \lambda(t) = \lambda(t) A \lambda(t)^{\mathsf{T}} = \begin{bmatrix} t^{2k}a_1 & 0 & t^{2k-2}a_2 & \cdots & a_{k+1} \\ 0 & -t^{2k-2}a_2 & 0 & \cdots & 0 \\ t^{2k-2}a_2 & 0 & t^{2k-4}a_3 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{k+1} & 0 & 0 & \cdots & 0 \end{bmatrix}.$$

It follows immediately that

$$\lim_{t\to 0} A \cdot \lambda(t) = A_{(n)},$$

where $A_{(n)}$ is given by (3.8).

□

Recall the following well-known theorem of Bialyncki-Birula.

Theorem 5.2. *[1, Theorem 4.3] Let* Z *be a smooth, irreducible projective variety with a (right) action of a* 1*-dimensional torus* $T' \cong \mathbb{C}^*$ *such that the fixed point set is finite. Then*

1. *For each fixed point* $x \in Z^{T'}$*, the set*

   $$C_x := \{y \in Z : \lim_{t \to 0} y \cdot t = x\}$$

   is an affine space.

2. *The closures* $\overline{C_x}$ *of the cells* C_x *form an additive basis for the Chow ring (as well as the cohomology ring) of* Z*.*

We would like to be able to deduce statement 2 in Theorem 5.2 in our setting where the variety Y_n is neither smooth nor irreducible. In order to accomplish this, we use a simplified version of a result of Carrell and Goresky.

Theorem 5.3. *[3, Theorem 1] Let* Z *be a (possibly singular, possibly reducible) variety which admits a (right)* $\mathbb{C}^*$*-action whose fixed point set is finite. If statement 1 holds in Theorem 5.2, then so does statement 2.*

We call the decomposition

$$Z = \bigsqcup_{x \in Z^{T'}} C_x$$

the *BB-decomposition* of Z with respect to the one-parameter subgroup T'; the varieties C_x are the *cells* of Z.

Theorem 5.4. *The decomposition*

$$Y_n = \bigsqcup_{\gamma \in F_n} Y_{n,\gamma}$$

is a BB-decomposition of Y_n *with respect to* T' *given by (3.6). In particular, the cell closures* $\overline{Y_{n,\gamma}}$ *form an additive basis for* $H^*(Y_n)$*. Moreover, if* $\gamma \in F_n$ *is a composition with* k *parts,* $\dim Y_{n,\gamma} = (n-k)/2$.

Proof. Proposition 4.3 and Lemma 5.1 verify statement 1 in Theorem 5.2, so we may apply Theorem 5.3. □

6 The Poincaré Polynomial

Using Theorem 5.4 we compute the Poincaré polynomial of Y_n:

$$P_{Y_n}(q) = \sum_{\gamma\in F_n} q^{\dim Y_{n,\gamma}} = \sum_{\gamma=(\gamma_1,\ldots,\gamma_k)\in F_n} q^{(n-k)/2}.$$

Using (2.1) and (2.2) to partition F_n into $F'_n \sqcup F''_n$, it follows that

$$\begin{aligned} P_{Y_n}(q) &= \sum_{\gamma=(\gamma_1,\ldots,\gamma_k)\in F'_n} q^{(n-k)/2} &&+ \sum_{\rho=(\rho_1\ldots,\rho_l)\in F''_n} q^{(n-l)/2} \\ &= \sum_{\gamma=(\gamma_2,\ldots,\gamma_k)\in F_{n-1}} q^{[(n-1)-(k-1)]/2} &&+ \sum_{\rho=(\rho_1-2\ldots,\rho_l)\in F_{n-2}} q\cdot q^{[(n-2)-l]/2}, \end{aligned}$$

which simplifies to

$$P_{Y_n}(q) = P_{Y_{n-1}}(q) + qP_{Y_{n-2(q)}}.$$

It is easy to easy to check that $P_{Y_1}(q) = P_{Y_2}(q) = 1$.

Theorem 6.1. *The Poincaré polynomial of the unipotent fixed variety Y_n is given by*

$$P_{Y_n}(q) = \sum_{i=0}^{n} \binom{n-1-i}{i} q^i. \tag{6.2}$$

Remark 6.3. By the Hard Lefschetz Theorem, $P_{Y_n}(q)$ is unimodal.

Proof. Formula (6.2) follows from the basic recurrence

$$\binom{n-1-i}{i} = \binom{n-2-i}{i-1} + \binom{n-2-i}{i}$$

and induction. □

7 The Poset of Cell Closures

We put a natural geometric partial order on F_n.

Definition 7.1. For $\rho = (\rho_1, \ldots, \rho_l)$ and $\gamma = (\gamma_k, \ldots, \gamma_k)$ in F_n,

$$\rho \leq \gamma \iff Y_{n,\rho} \subseteq \overline{Y_{n,\gamma}}.$$

Proposition 7.2. *Let* $\gamma = (\gamma_1, \ldots, \gamma_k), \rho = (\rho_1, \ldots, \rho_l) \in F_n$. *Then,* γ *covers* ρ *if and only if there exists a part* γ_i *of* γ *which is* > 1 *and* ρ *is obtained from* γ *by replacing* γ_i *by* $\gamma_i - 2, 1, 1$, *i.e.* $\rho = (\gamma_1, \ldots, \gamma_{i-1}, \gamma_i - 2, 1, 1, \gamma_{i+1}, \ldots, \gamma_k)$. *Moreover,*

$$\overline{Y_{n,\gamma}} = \mathbb{P}^{(\gamma_1-1)/2} \times \cdots \times \mathbb{P}^{(\gamma_k-1)/2}.$$

Proof. This follows from the following explicit description of the the cell $Y_{n,\gamma}$:

$$Y_{n,\gamma} = \prod_{i=1}^{k} \mathcal{A}_{\gamma_i},$$

where

$$\mathcal{A}_m = \left\{ \begin{bmatrix} a_1 & 0 & a_2 & \cdots & a_{(m-1)/2} \\ 0 & -a_2 & 0 & \cdots & 0 \\ a_2 & 0 & a_3 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{(m-1)/2} & 0 & 0 & \cdots & 0 \end{bmatrix} : a_i \in \mathbb{K}, a_{(m-1)/2} \neq 0 \right\}$$

for m odd. Note that $\mathcal{A}_m \cong \mathbb{A}^{(m-1)/2}$.

It is easy to see that $\overline{\mathcal{A}_m} = \mathcal{A}_m \cup \mathcal{A}_{m-2} \cup \cdots \cup \mathcal{A}_1$, the standard decomposition of $\mathbb{P}^{(m-1)/2}$. □

Corollary 7.3. *Let* $\gamma = (\gamma_1, \gamma_2, \ldots, \gamma_k) \in F_n$ *and let* $\hat{0} = (1, 1, \ldots, 1)$, *the minimal element of* F_n. *The interval* $[\hat{0}, \gamma]$ *is a product of chains. More specifically,*

$$[\hat{0}, \gamma] = C_{(\gamma_1-1)/2} \times C_{(\gamma_2-1)/2} \times \cdots \times C_{(\gamma_k-1)/2},$$

where C_m *denotes the chain with* $m + 1$ *elements.*

Note that irreducible components of Y_n correspond to maximal elements of $(F_n, \leq)$. Also, F_n has a $\hat{0}$, namely the composition $(1, 1, \ldots, 1)$ consisting of n 1's; however, F_n has more than one maximal element when $n \geq 4$.

Example 7.4. For $n = 6$, we have the following Hasse diagram, graded by dimension (see Lemma 7.6):

We conclude from Figure 7.1 that Y_6 has four irreducible components: two of which are isomorphic to $\mathbb{P}^2$, one isomorphic to $\mathbb{P}^1 \times \mathbb{P}^1$ and one isomorphic to $\mathbb{P}^1$.

Remark 7.5. It is clear from Figure 7.1 that unlike the case of Springer fibers, the irreducible components of Y_n are not equidimensional.



Figure 7.1: Hasse diagram for Y_6.

Lemma 7.6. *The poset* $(F_n, \leq)$ *is a graded poset. For* $\gamma = (\gamma_1, \gamma_2, \ldots, \gamma_k) \in F_n$, *the rank function is given by* $\ell(\gamma) = (n-k)/2 = \dim Y_{n,\gamma}$.

Proof. Clearly, if $\rho \leq \gamma$, then $\ell(\rho) = \dim Y_{n,\rho} \leq \dim Y_{n,\gamma} = \ell(\gamma)$. Also, if γ covers ρ, then by Proposition 7.2, ρ is obtained from γ by replacing some γ_i by $\gamma_i - 2, 1, 1$. Thus,

$$\ell(\rho) = (n - (k+2))/2 = \ell(\gamma) - 1.$$

□

Lemma 7.7. *Let* $\gamma = (\gamma_1, \ldots, \gamma_k)$ *and* $\tau = (\tau_1, \ldots, \tau_l)$ *be two elements from* F_n *such that* $\gamma_1 > 1$ *and* $\tau_1 = 1$. *If* $\tau \leq \gamma$, *then* $\tau_1 = \cdots = \tau_{\gamma_1} = 1$.

Proof. Since $\gamma \geq \tau$, there exists a sequence of coverings $\gamma = \gamma^{(0)} \geq \gamma^{(1)} \geq \cdots \geq \gamma^{(s)} = \tau$ and a subsequence $\gamma = \gamma^{(i_0)} \geq \gamma^{(i_1)} \geq \cdots \geq \gamma^{(i_d)} = \tau$ such that

$$\begin{gathered}
\gamma_1^{(i_1)} = \gamma_1 - 2 \text{ and } \gamma_2^{(i_1)} = \gamma_3^{(i_1)} = 1, \\
\gamma_1^{(i_2)} = \gamma_1^{(i_1)} - 2 \text{ and } \gamma_2^{(i_2)} = \gamma_3^{(i_2)} = \gamma_4^{(i_2)} = \gamma_5^{(i_2)} = 1, \\
\vdots \\
\gamma_1^{(i_d)} = \gamma_1^{(i_{d-1})} - 2 = 1 \text{ and } \gamma_2^{(i_d)} = \cdots = \gamma_{2d+1}^{(i_d)} = 1.
\end{gathered}$$

It follows that $\tau_1 = \cdots = \tau_j = 1$ for some $j \geq 2d + 1 = \gamma_1$.

□

Theorem 7.8. *The Fibonacci poset* $(F_n, \leq)$ *is a meet-semi lattice.*

Proof. The proof is by induction on n. Assume that the statement is true for F_m with $m < n$.

Recall the decomposition $F_n = F'_n \sqcup F''_n$ and the bijections ϕ (2.1) and ψ (2.2). Let γ and ρ be two elements from F_n. If γ and ρ both lie in F'_n or both lie in F''_n, then by the induction hypothesis there exists a unique $\gamma \wedge \rho$. Therefore, without loss of generality, we may assume that $\gamma \in F''_n$ and $\rho \in F'_n$.

Let $\widetilde{\gamma} \in F_n$ be the element obtained from γ by replacing γ_1 by the sequence $1, \ldots, 1$ of γ_1 1's. Then $\widetilde{\gamma} \in F'_n$, and by the induction hypothesis $\widetilde{\gamma} \wedge \rho \in F'_n$ exists uniquely. If $\tau \leq \gamma$ and $\tau \leq \rho$, then by Lemma 7.7, $\tau \leq \widetilde{\gamma}$, so $\tau \leq \widetilde{\gamma} \wedge \rho$.

□

Theorem 7.9. *Let a_n be the number of irreducible components of Y_n. Then*

$$a_n = a_{n-1} + a_{n-3} \qquad \text{for } n \geq 4.$$

The initial values of the sequence are $a_1 = a_2 = a_3 = 1$.

Proof. Let $F_n^{\max} \subseteq F_n$ denote the set of maximal elements in F_n. From Proposition 7.2, a composition $\gamma \in F_n$ belongs to $F_n^{\max}$ if and only if the composition contains no consecutive 1's, except possibly two consecutive 1's at the beginning.

Let A, respectively A', be the subsets of $F_n^{\max}$ consisting of elements whose last part is 1, respectively ≥ 5. Let $B \subset F_n^{\max}$ be the subset consisting of elements whose last part is 3. F_n is a disjoint union of A, A', and B.

Define $f : A \cup A' \to F_{n-1}^{\max}$ by sending $\gamma = (\gamma_1, \ldots, \gamma_k)$ to $(\gamma_1, \ldots, \gamma_{k-1})$ if γ is in A, and to $(\gamma_1, \ldots, \gamma_k - 2, 1)$ if γ is in A'. Similarly, define $g : B \to F_{n-3}^{max}$ by $g(\gamma) = (\gamma_1, \ldots, \gamma_{k-1})$. The maps f and g are bijections, and therefore,

$$a_n = |F_n^{\max}| = |F_{n-1}^{\max}| + |F_{n-3}^{\max}| = a_{n-1} + a_{n-3}.$$

□

Remark 7.10. The (ordinary) generating function for $\{a_n\}$ (with the convention $a_0 = 1$) is

$$\sum_{n=0}^{\infty} a_n x^n = \frac{1}{1 - x - x^3}.$$

This sequence also counts the number of compositions of $n - 2$ into parts 1 and 2 with no consecutive 2's.

References

[1] A. Białynicki-Birula. Some theorems on actions of algebraic groups. *Ann. of Math. (2)*, 98:480–497, 1973.

[2] M.B. Can and M. Joyce. Unipotent invariant quadrics. Preprint. `http://arxiv.org/abs/1010.2187`, 2010.

[3] J. B. Carrell and R. M. Goresky. A decomposition theorem for the integral homology of a variety. *Invent. Math.*, 73(3):367–381, 1983.

[4] M. Chasles. Détermination du hombre des sections coniques qui doivent toucher cinq courbes données d'ordre quelconque, ou satisfaire à diverses autres conditions. *C. R. Acad. Sci. Paris*, 58:222–226, 1864.

[5] C. De Concini, M. Goresky, R. MacPherson, and C. Procesi. On the geometry of quadrics and their degenerations. *Comment. Math. Helv.*, 63(3):337–413, 1988.

[6] C. De Concini and C. Procesi. Complete symmetric varieties. In *Invariant theory (Montecatini, 1982)*, volume 996 of *Lecture Notes in Math.*, pages 1–44. Springer, Berlin, 1983.

[7] C. De Concini and T. A. Springer. Betti numbers of complete symmetric varieties. In *Geometry today (Rome, 1984)*, volume 60 of *Progr. Math.*, pages 87–107. Birkhäuser Boston, Boston, MA, 1985.

[8] Hermann Schubert. *Kalkül der abzählenden Geometrie*. Springer-Verlag, Berlin, 1979. Reprint of the 1879 original, With an introduction by Steven L. Kleiman.

[9] Nicolas Spaltenstein. On the fixed point set of a unipotent element on the variety of Borel subgroups. *Topology*, 16(2):203–204, 1977.

[10] T. A. Springer. Trigonometric sums, Green functions of finite groups and representations of Weyl groups. *Invent. Math.*, 36:173–207, 1976.

[11] T. A. Springer. A construction of representations of Weyl groups. *Invent. Math.*, 44(3):279–293, 1978.

[12] Richard P. Stanley. *Enumerative combinatorics. Vol. 1*, volume 49 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1997. With a foreword by Gian-Carlo Rota, Corrected reprint of the 1986 original.

[13] Elisabetta Strickland. Schubert-type cells for complete quadrics. *Adv. in Math.*, 62(3):238–248, 1986.